

# Instantons and Chern-Simons Flows in 6, 7 and 8 dimensions

Olaf Lechtenfeld

*Institut für Theoretische Physik, Leibniz Universität Hannover  
Appelstraße 2, 30167 Hannover, Germany  
URL: [www.itp.uni-hannover.de/~lechtenf](http://www.itp.uni-hannover.de/~lechtenf)*

and

*Centre for Quantum Engineering and Space-Time Research  
Leibniz Universität Hannover  
Welfengarten 1, 30167 Hannover, Germany*

## Abstract

The existence of  $K$ -instantons on a cylinder  $M^7 = \mathbb{R}_\tau \times \frac{K}{H}$  over a homogeneous nearly Kähler 6-manifold  $\frac{K}{H}$  requires a conformally parallel or a cocalibrated  $G_2$ -structure on  $M^7$ . The generalized anti-self-duality on  $M^7$  implies a Chern-Simons flow on  $\frac{K}{H}$  which runs between instantons on the coset. For  $K$ -equivariant connections, the torsionful Yang-Mills equation reduces to a particular quartic dynamics for a Newtonian particle on  $\mathbb{C}$ . When the torsion corresponds to one of the  $G_2$ -structures, this dynamics follows from a gradient or hamiltonian flow equation, respectively. We present the analytic (kink-type) solutions and plot numerical non-BPS solutions for general torsion values interpolating between the instantonic ones.

# 1 Introduction

Yang-Mills instantons exist dimensions  $d$  larger than four only when there is additional geometric structure on the manifold  $M^d$  (besides the Riemannian one). In order to formulate generalized first-order anti-self-duality conditions which imply the second-order Yang-Mills equations (possibly with torsion),  $M^d$  must be equipped with a so called  $G$ -structure, which is a globally defined but not necessarily closed  $(d-4)$ -form  $\Sigma$ , so that the weak holonomy group of  $M^d$  gets reduced.

Instanton solutions in higher dimensions are rare in the literature. In the mid-eighties, Fairlie and Nuyts and also Fubini and Nicolai discovered the Spin(7)-instanton on  $\mathbb{R}^8$ . Eight years later, a similar  $G_2$ -instanton on  $\mathbb{R}^7$  was found by Ivanova and Popov and also by Günaydin and Nicolai. Our recent work shows that these so called octonionic instantons are not isolated but embedded into a whole family living on a class of conical non-compact manifolds [1].

The string vacua in heterotic flux compactifications contain non-abelian gauge fields which in the supergravity limit are subject to Yang-Mills equations with torsion  $\mathcal{H}$  determined by the three-form flux. Prominent cases admitting instantons are  $\text{AdS}_{10-d} \times M^d$ , where  $M^d$  is equipped with a  $G$ -structure, with  $G$  being SU(3),  $G_2$  or Spin(7) for  $d = 6, 7$  or 8, respectively. Homogeneous nearly Kähler 6-manifolds  $\frac{K}{H}$  and (iterated) cylinders and (sine-)cones over them provide simple examples, for which all  $K$ -equivariant Yang-Mills connections can be constructed [2, 3]. Natural choices for the gauge group are  $K$  or  $G$ .

Clearly, the Yang-Mills instantons discussed here serve to construct heterotic string solitons, as was first done in 1990 by Strominger for the gauge five-brane. It is therefore of interest to extend our new instantons to solutions of (string-corrected) heterotic supergravity and obtain novel string/brane vacua [4, 5, 6].

In this talk, I present the construction for the simplest case of a cylinder over a compact homogeneous nearly Kähler coset  $\frac{K}{H}$ , which allows for a conformally parallel or a cocalibrated  $G_2$ -structure. I display a family of non-BPS Yang-Mills connections, which contain two instantons at distinguished parameter values corresponding to those  $G_2$ -structures. In these two cases, anti-self-duality implies a Chern-Simons flow on  $\frac{K}{H}$ .

Finally, I must apologize for the omission – due to page limitation – of all relevant literature besides my own papers on which this talk is based. The reader can find all references therein.

## 2 Self-duality in higher dimensions

The familiar four-dimensional anti-self-duality condition for Yang-Mills fields  $F$  may be generalized to suitable  $d$ -dimensional Riemannian manifolds  $M$ ,

$$*F = -\Sigma \wedge F \quad \text{for} \quad F = dA + A \wedge A \quad \text{and} \quad \Sigma \in \Lambda^{d-4}(M), \quad (1)$$

if there exists a geometrically natural  $(d-4)$ -form  $\Sigma$  on  $M$ . Applying the gauge-covariant derivative  $D = d + [A, \cdot]$  it follows that

$$D*F + d\Sigma \wedge F = 0 \quad \Longleftrightarrow \quad \text{Yang-Mills with torsion} \quad \mathcal{H} = *d\Sigma \in \Lambda^3(M). \quad (2)$$

This torsionful Yang-Mills equation extremizes the action

$$S_{\text{YM}} + S_{\text{CS}} = \int_M \text{tr} \{ F \wedge *F + (-)^{d-3} \Sigma \wedge F \wedge F \} = \int_M \text{tr} \{ F \wedge *F + \frac{1}{2} d\Sigma \wedge (A dA + \frac{2}{3} A^3) \}. \quad (3)$$

Related to this generalized anti-self-duality is the gradient Chern-Simons flow on  $M$ ,


$$\frac{dA}{d\tau} = \frac{\delta}{\delta A} S_{\text{CS}} = *(\text{d}\Sigma \wedge F) \sim *\text{d}\Sigma \lrcorner F. \quad (4)$$

In fact, this equation follows from generalized anti-self-duality on the cylinder  $\widetilde{M} = \mathbb{R}_\tau \times M$  over  $M$  (in the  $A_\tau=0$  gauge).

The question is therefore: Which manifolds admit a global  $(d-4)$ -form? And the answer is:  $G$ -structure manifolds, i.e. manifolds with a weak special holonomy. Let me give the key cases we shall encounter in this talk:

| $d$ | $G$     | $\Sigma$ | cases                 | example                                                  | structure                                                                                   |
|-----|---------|----------|-----------------------|----------------------------------------------------------|---------------------------------------------------------------------------------------------|
| 6   | SU(3)   | $\omega$ | Kähler                | $\mathbb{C}P^3$                                          | $\text{d}\omega = 0$                                                                        |
| 6   | SU(3)   | $\omega$ | nearly Kähler         | $S^6 = \frac{G_2}{\text{SU}(3)}$                         | $\text{d}\omega \sim \text{Im}\Omega, \text{dRe}\Omega \sim \omega^2$                       |
| 7   | $G_2$   | $\psi$   | conf. parallel $G_2$  | $\mathbb{R}_\tau \times$ nearly Kähler                   | $\text{d}\psi \sim \psi \wedge \text{d}\tau, \text{d}*\psi \sim -*\psi \wedge \text{d}\tau$ |
| 7   | $G_2$   | $\psi$   | nearly parallel $G_2$ | $X_{k,\ell} = \frac{\text{SU}(3)}{\text{U}(1)_{k,\ell}}$ | $\text{d}\psi \sim *\psi \Rightarrow \text{d}*\psi = 0$                                     |
| 7   | $G_2$   | $\psi$   | parallel $G_2$        | cone(nearly Kähler)                                      | $\text{d}\psi = 0 = \text{d}*\psi$                                                          |
| 8   | Spin(7) | $\Sigma$ | parallel Spin(7)      | $\mathbb{R}_\tau \times$ parallel $G_2$                  | $\text{d}\Sigma = 0, *\Sigma = \Sigma$                                                      |

Some of those cases are related via the following scheme, with examples in square brackets.



For this talk I shall consider (reductive non-symmetric) coset spaces  $M = \frac{K}{H}$  in  $d=6$  as well as cylinders and cones over them. In all these cases, the gauge group is chosen to be  $K$ .

### 3 Six dimensions: nearly Kähler coset spaces

All known compact nearly Kähler 6-manifolds  $M^6$  are nonsymmetric coset spaces  $K/H$ :

$$S^6 = \frac{G_2}{\text{SU}(3)}, \quad \frac{\text{Sp}(2)}{\text{Sp}(1) \times \text{U}(1)}, \quad \frac{\text{SU}(3)}{\text{U}(1) \times \text{U}(1)}, \quad S^3 \times S^3 = \frac{\text{SU}(2) \times \text{SU}(2) \times \text{SU}(2)}{\text{SU}(2)}. \quad (5)$$

The coset structure  $H \triangleleft K$  implies the decomposition

$$\text{Lie}(K) \equiv \mathfrak{k} = \mathfrak{h} \oplus \mathfrak{m} \quad \text{with} \quad \mathfrak{h} \equiv \text{Lie}(H) \quad \text{and} \quad [\mathfrak{h}, \mathfrak{m}] \subset \mathfrak{m}. \quad (6)$$

Interestingly, the reflection automorphism of symmetric spaces gets generalized to a so called tri-symmetry automorphism  $S : K \rightarrow K$  with  $S^3 = \text{id}$  implying

$$s : \mathfrak{k} \rightarrow \mathfrak{k} \quad \text{with} \quad s|_{\mathfrak{h}} = \mathbb{1} \quad \text{and} \quad s|_{\mathfrak{m}} = -\frac{1}{2} + \frac{\sqrt{3}}{2} J = \exp\{\frac{2\pi}{3} J\} , \quad (7)$$

effecting a  $\frac{2\pi}{3}$  rotation on  $TM^6$ . I pick a Lie-algebra basis

$$\{I_{a=1,\dots,6}, I_{i=7,\dots,\dim G}\} \quad \text{with} \quad [I_a, I_b] = f_{ab}^i I_i + f_{ab}^c I_c , \quad (8)$$

involving the structure constants  $f_{ab}^\bullet$ . The Cartan-Killing form then reads

$$\langle \cdot, \cdot \rangle_{\mathfrak{k}} = -\text{tr}_{\mathfrak{k}}(\text{ad}(\cdot) \circ \text{ad}(\cdot)) = 3 \langle \cdot, \cdot \rangle_{\mathfrak{h}} = 3 \langle \cdot, \cdot \rangle_{\mathfrak{m}} = \mathbb{1} . \quad (9)$$

Expanding all structures in a basis of canonical one-forms  $e^a$  framing  $T^*(G/H)$ ,

$$g = \delta_{ab} e^a e^b , \quad \omega = \frac{1}{2} J_{ab} e^a \wedge e^b , \quad \Omega = -\frac{1}{\sqrt{3}} (f + iJf)_{abc} e^a \wedge e^b \wedge e^c , \quad (10)$$

we see that the almost complex structure  $(J_{ab})$  and the structure constants  $f_{abc}$  rule everything.

Nearly Kähler 6-manifolds are special in that the torsion term in (2) *vanishes by itself*! What is more, this property is actually *equivalent* to the generalized anti-self-duality condition (1):

$$*F = -\omega \wedge F \quad \Longleftrightarrow \quad 0 = d\omega \wedge F \sim \text{Im}\Omega \wedge F \quad \Longleftrightarrow \quad \text{DUY equations} , \quad (11)$$

where the Donaldson-Uhlenbeck-Yau (DUY) equations<sup>1</sup> state that

$$F^{2,0} = F^{0,2} = 0 \quad \text{and} \quad \omega \lrcorner F = 0 . \quad (12)$$

Another interpretation of this anti-self-duality condition is that it projects  $F$  to the 8-dimensional eigenspace of the endomorphism  $*(\omega \wedge \cdot)$  with eigenvalue  $-1$ , which contains the part of  $F^{1,1}$  orthogonal to  $\omega$ . The equations (11) imply also  $\text{Re}\Omega \wedge F = 0$  and the (torsion-free) Yang-Mills equations  $D*F = 0$ . Clearly, they separately extremize both  $S_{\text{YM}}$  and  $S_{\text{CS}}$  in (3), but of course yield only BPS-type classical solutions. In components the above relations take the form

$$\frac{1}{2} \epsilon_{abcdef} F_{ef} = -J_{[ab} F_{cd]} \quad \Longleftrightarrow \quad 0 = f_{abc} F_{bc} , \quad (13)$$

$$\implies \quad \omega_{ab} F_{ab} = 0 , \quad (Jf)_{abc} F_{bc} = 0 , \quad D_a F_{ab} = 0 . \quad (14)$$

I notice that each Chern-Simons flow  $\dot{A}_a \sim f_{abc} F_{bc}$  on  $M^6$  ends in an instanton.

Let me look for  $K$ -equivariant connections  $A$  on  $M^6$ . If I restrict their value to  $\mathfrak{h}$ , the answer is unique: the only ‘ $H$ -instanton’ is the so called canonical connection

$$A^{\text{can}} = e^i I_i \quad \longrightarrow \quad F^{\text{can}} = -\frac{1}{2} f_{ab}^i e^a \wedge e^b I_i , \quad (15)$$

where  $e^i = e_a^i e^a$ . Generalizing to ‘ $K$ -instantons’, I extend to

$$A = e^i I_i + e^a \Phi_{ab} I_b \quad \text{with ansatz} \quad (\Phi_{ab}) =: \Phi = \phi_1 \mathbb{1} + \phi_2 J , \quad (16)$$

which is in fact general for  $G_2$  invariance on  $S^6$ . Its curvature is readily computed to

$$F_{ab} = F_{ab}^{1,1} + F_{ab}^{2,0 \oplus 0,2} = (|\Phi|^2 - 1) f_{ab}^i I_i + [(\bar{\Phi}^2 - \Phi) f]_{abc} I_c \quad (17)$$

and displays the tri-symmetry invariance under  $\Phi \rightarrow \exp\{-\frac{2\pi}{3} J\} \Phi$ . The solutions to the BPS conditions (11) are

$$\bar{\Phi}^2 = \Phi \quad \implies \quad \Phi = 0 \quad \text{or} \quad \Phi = \exp\{\frac{2\pi k}{3} J\} \quad \text{for} \quad k = 0, 1, 2 , \quad (18)$$

which yields three flat  $K$ -instanton connections besides the canonical curved one,

$$A^{(k)} = e^i I_i + e^a (s^k I)_a \quad \text{and} \quad A^{\text{can}} = e^i I_i . \quad (19)$$

---

<sup>1</sup>also known as ‘hermitian Yang-Mills equations’

## 4 Seven dimensions: cylinder over nearly Kähler cosets

Let me step up one dimension and consider 7-manifolds  $M^7$  with weak  $G_2$  holonomy associated with a  $G_2$ -structure three-form  $\psi$ . Here, the 7 generalized anti-self-duality equations project  $F$  onto the  $-1$  eigenspace of  $\ast(\psi \wedge \cdot)$ , which is 14-dimensional and isomorphic to the Lie algebra of  $G_2$ ,

$$\ast F = -\psi \wedge F \iff \ast \psi \wedge F = 0 \iff \psi \lrcorner F = 0, \quad (20)$$

providing an alternative form of the condition. In components, it reads

$$\frac{1}{2}\epsilon_{abcdefg}F_{fg} = -\psi_{[abc}F_{de]} \iff 0 = \psi_{abc}F_{bc}. \quad (21)$$

For the parallel and nearly parallel  $G_2$  cases, the previous accident (11) recurs,

$$d\psi \sim \ast \psi \implies d\psi \wedge F = 0 \implies D \ast F = 0, \quad (22)$$

and the torsion decouples. Note that on a general weak  $G_2$ -manifold there are two different flows,

$$\frac{dA(\sigma)}{d\sigma} = \ast d\psi \lrcorner F(\sigma) \quad \text{and} \quad \frac{dA(\sigma)}{d\sigma} = \psi \lrcorner F(\sigma) \quad \text{for } \sigma \in \mathbb{R}, \quad (23)$$

which coincide in the nearly parallel case. The second flow ends in an instanton on  $M^7$ .

In this talk I focus on cylinders  $M^7 = \mathbb{R}_\tau \times \frac{K}{H}$  over nearly Kähler cosets, with a metric  $g = (d\tau)^2 + \delta_{ab} e^a e^b$ , on which I study the Yang-Mills equation with a torsion given by

$$\ast \mathcal{H} = \frac{1}{3}\kappa d\omega \wedge d\tau \iff T_{abc} = \kappa f_{abc} \quad (24)$$

with a real parameter  $\kappa$ . We shall see that for special values of  $\kappa$  my torsionful Yang-Mills equation

$$D \ast F + \frac{1}{3}\kappa d\omega \wedge d\tau \wedge F = 0 \quad (25)$$

descends from an anti-self-duality condition (20).

Taking the  $A_0=0$  gauge and borrowing the ansatz (16) from the nearly Kähler base, I write

$$A_a = e_a^i I_i + [\Phi(\tau) I]_a \implies F_{0a} = [\dot{\Phi} I]_a, \quad F_{ab} = (|\Phi|^2 - 1) f_{ab}^i I_i + [(\bar{\Phi}^2 - \Phi) f]_{abc} I_c \quad (26)$$

which depends on a complex function  $\Phi(\tau)$  (values in the  $(\mathbb{1}, J)$  plane). Sticking this into (25) and computing for a while, one arrives at

$$\ddot{\Phi} = (\kappa - 1)\Phi - (\kappa + 3)\bar{\Phi}^2 + 4\bar{\Phi}\Phi^2 =: \frac{1}{3}\frac{\partial V}{\partial \Phi}. \quad (27)$$

Nice enough, I have obtained a  $\phi^4$  model with an action

$$S[\Phi] \sim \int_{\mathbb{R}} d\tau \{3|\dot{\Phi}|^2 + V(\Phi)\} \quad \text{for} \quad V(\Phi) = (3 - \kappa) + 3(\kappa - 1)|\Phi|^2 - (3 + \kappa)(\Phi^3 + \bar{\Phi}^3) + 6|\Phi|^4 \quad (28)$$

devoid of rotational symmetry (for  $\kappa \neq -3$ ) but enjoying tri-symmetry in the complex plane. It leads me to a mechanical analog problem of a Newtonian particle on  $\mathbb{C}$  in a potential  $-V$ . I obtain the same action by plugging (26) directly into (3) with  $d\Sigma = \ast \mathcal{H}$  from (24).

For the case of  $\frac{K}{H} = S^6 = \frac{G_2}{\text{SU}(3)}$ , equation (27) produces in fact *all*  $G$ -equivariant Yang-Mills connections on  $\mathbb{R}_\tau \times \frac{K}{H}$ . On  $\frac{\text{Sp}(2)}{\text{Sp}(1) \times \text{U}(1)}$  and  $\frac{\text{SU}(3)}{\text{U}(1) \times \text{U}(1)}$ , however, the most general  $G$ -equivariant connections involves two respective three complex functions of  $\tau$ . The corresponding Newtonian dynamics on  $\mathbb{C}^2$  respective  $\mathbb{C}^3$  is of similar type but constrained by the conservation of Noether charges related to relative phase rotations of the complex functions.

## 5 Seven dimensions: solutions

Finite-action solutions require Newtonian trajectories between zero-potential critical points  $\hat{\Phi}$ . With two exotic exceptions,  $dV(\hat{\Phi}) = 0 = V(\hat{\Phi})$  yields precisely the BPS configurations on  $\frac{K}{H}$ :

- $\hat{\Phi} = e^{2\pi i k/3}$  with  $V(\hat{\Phi}) = 0$  for all values of  $\kappa$  and  $k = 0, 1, 2$
- $\hat{\Phi} = 0$  with  $V(\hat{\Phi}) = 3 - \kappa$  vanishing only at  $\kappa = 3$

Kink solutions will interpolate between two different critical points, while bounces will return to the critical starting point. Thus for generic  $\kappa$  values one may have kinks of ‘transversal’ type, connecting two third roots of unity, as well as bounces. For  $\kappa = 3$  ‘radial’ kinks, reaching such a root from the origin, may occur as well. Numerical analysis reveals the domains of existence in  $\kappa$ :

| $\kappa$ interval   | $(-\infty, -3]$  | $(-3, +3)$          | $+3$           | $(+3, +5)$       | $[+5, +\infty)$ |
|---------------------|------------------|---------------------|----------------|------------------|-----------------|
| types of trajectory | radial<br>bounce | transversal<br>kink | radial<br>kink | radial<br>bounce | –<br>–          |

On the following page I display contour plots of the potential and finite-action trajectories for eight choices of  $\kappa$ . They reveal three special values of  $\kappa$ : At  $\kappa = -3$  rotational symmetry emerges; this is a degenerate situation. At  $\kappa = -1$  and at  $\kappa = +3$ , the trajectories are straight, indicating integrability. Indeed, behind each of these two cases lurks a first-order flow equation, which originates from anti-self-duality and hence a particular  $G_2$ -structure  $\psi$ .

Let me first discuss  $\kappa = +3$ . For this value I find that

$$3\ddot{\Phi} = \frac{\partial V}{\partial \Phi} \iff \sqrt{2}\dot{\Phi} = \pm \frac{\partial W}{\partial \Phi} \quad \text{with} \quad W = \frac{1}{3}(\Phi^3 + \bar{\Phi}^3) - |\Phi|^2, \quad (29)$$

which is a gradient flow with a real superpotential  $W$ , as

$$V = 6 \left| \frac{\partial W}{\partial \Phi} \right|^2 \quad \text{for} \quad \kappa = +3. \quad (30)$$

It admits the obvious analytic radial kink solution,

$$\Phi(\tau) = e^{\frac{2\pi i k}{3}} \left( \frac{1}{2} \pm \frac{1}{2} \tanh \frac{\tau}{2\sqrt{3}} \right). \quad (31)$$

What is the interpretation of this gradient flow in terms of the original Yang-Mills theory? Demanding that the torsion in (24) comes from a  $G_2$ -structure,  $*\mathcal{H} = d\psi$ , I am led to

$$\psi = \frac{1}{3}\kappa \omega \wedge d\tau + \alpha \text{Im}\Omega \implies d\psi \sim \kappa \text{Im}\Omega \wedge d\tau \sim \psi \wedge d\tau \quad (32)$$

where  $\alpha$  is undetermined. This is a conformally parallel  $G_2$ -structure, and (20) quantizes the coefficients to  $\alpha = 1$  and  $\kappa = 3$ , fixing

$$\psi = \omega \wedge d\tau + \text{Im}\Omega = r^{-3}(r^2\omega \wedge dr + r^3\text{Im}\Omega) = r^{-3}\psi_{\text{cone}} \quad \text{with} \quad e^\tau = r, \quad (33)$$









where I displayed the conformal relation to the parallel  $G_2$ -structure on the cone over  $\frac{K}{H}$ .

Alternatively, with this  $G_2$ -structure the 7 anti-self-duality equations (20) turn into

$$\omega \lrcorner F \sim J_{ab} F_{ab} = 0 \quad \text{and} \quad \dot{A} \sim d\omega \lrcorner F \sim e^a f_{abc} F_{bc}. \quad (34)$$

With the ansatz (26), the first relation is automatic, and the second one indeed reduces to (29). As a consistency check, one may verify that

$$\int_{\frac{K}{H}} \text{tr} \{ \omega \wedge F \wedge F \} \propto W(\Phi) + \frac{1}{3}. \quad (35)$$



I now come to the other instance of straight trajectories,  $\kappa=-1$ . For this value I find that

$$3\ddot{\Phi} = \frac{\partial V}{\partial \Phi} \quad \Longleftarrow \quad \sqrt{2}\dot{\Phi} = \pm i \frac{\partial H}{\partial \Phi} \quad \text{with} \quad H = \frac{1}{3}(\Phi^3 + \bar{\Phi}^3) - |\Phi|^2, \quad (36)$$

which is a hamiltonian flow (note the imaginary multiplier!), running along the level curves of the function  $H$ , that is identical to  $W$ . It has the obvious analytic transverse kink solution,

$$\Phi(\tau) = -\frac{1}{2} \pm \frac{\sqrt{3}}{2} i (\tanh \frac{\tau}{2}) \quad (37)$$

and its images under the tri-symmetry action.

Have I discovered another hidden  $G_2$ -structure here? Let me try the other obvious choice,

$$\tilde{\psi} = \frac{1}{3}\tilde{\kappa}\omega \wedge d\tau + \tilde{\alpha} \text{Re}\Omega \quad \Longrightarrow \quad d\tilde{\psi} \sim \tilde{\kappa} \text{Im}\Omega \wedge d\tau + 2\tilde{\alpha}\omega \wedge \omega, \quad (38)$$

with coefficients  $\tilde{\kappa}$  and  $\tilde{\alpha}$  to be determined. It has not appeared in my table in Section 2, but obeys  $d * \tilde{\psi} = 0$ , which is known as a *cocalibrated*  $G_2$ -structure. But can it produce the proper torsion,

$$d\tilde{\psi} \wedge F \sim (\tilde{\kappa} \text{Im}\Omega \wedge d\tau + 2\tilde{\alpha}\omega \wedge \omega) \wedge F \stackrel{!}{=} -\text{Im}\Omega \wedge d\tau \wedge F ? \quad (39)$$

Employing the anti-self-duality with respect to  $\tilde{\psi}$ ,

$$*\tilde{\psi} \wedge F = 0 \quad \Longrightarrow \quad \omega \wedge \omega \wedge F = 2 \text{Im}\Omega \wedge d\tau \wedge F, \quad (40)$$

it works out, adjusting the coefficients to  $\tilde{\kappa}=3$  and  $\tilde{\alpha}=-1$ . Hence, my cocalibrated  $G_2$ -structure



$$\tilde{\psi} = \omega \wedge d\tau - \text{Re}\Omega \quad (41)$$

is responsible for the hamiltonian flow.

To see this directly, I import (41) into (20) and get

$$J_{ab} F_{ab} = 0 \quad \text{and} \quad \dot{A}_a \sim [Jf]_{abc} F_{bc}. \quad (42)$$

Again, the ansatz (26) fulfills the first relation, but the second one nicely turns into (36).




The story has an eight-dimensional twist, which can be inferred from the diagram in Section 2. There it is indicated that my cylinder is embedded into an 8-manifold  $M^8$  equipped with a parallel Spin(7)-structure  $\Sigma$ . It can be regarded as the cylinder over the cone over  $\frac{K}{H}$ . The four-form  $\Sigma$  descends to the cocalibrated  $G_2$ -structure  $\tilde{\psi}$ , while  $\psi$  is obtained by reducing to the cone and applying a conformal transformation.

The anti-self-duality condition on  $M^8$  represents 7 relations, which project  $F_8$  to the 21-dimensional  $-1$  eigenspace of  $*(\Sigma \wedge \cdot)$ . Contrary to the  $G_2$  situation (34), where 7 anti-self-duality equations split to 6 flow equations and the supplementary condition  $\omega \lrcorner F = 0$ , for Spin(7) the count precisely matches, as I have also 7 flow equations. Indeed, there is equivalence:

$$*_8 F_8 = -\Sigma \wedge F_8 \quad \Longleftrightarrow \quad \frac{\partial A_7(\sigma)}{\partial \sigma} = *_7(d\psi \wedge F_7(\sigma)) . \quad (43)$$

## 6 Partial summary

Let me schematically sum up the construction.



are  $G_2$ -instantons for Yang-Mills with torsion  $D * F + (*\mathcal{H}) \wedge F = 0$

from  $S[A] = \int_{\mathbb{R} \times \frac{K}{H}} \text{tr} \{ F \wedge *F + \frac{1}{3} \kappa \omega \wedge d\tau \wedge F \wedge F \}$  with  $\kappa = +3$  or  $-1$

and obey gradient/hamiltonian flow equations for  $\int_{\frac{K}{H}} \text{tr} \{ \omega \wedge F \wedge F \} \propto W(\Phi) + \frac{1}{3}$ .

## References

- [1] Gemmer K.P., Lechtenfeld O., Nölle C., Popov A.D. // JHEP 2011. V.09. P.103; arXiv:1108.3951.
- [2] Harland D., Ivanova T.A., Lechtenfeld O., Popov A.D. // Commun. Math. Phys. 2010. V.300. P.185; arXiv:0909.2730.
- [3] Bauer I., Ivanova T.A., Lechtenfeld O., Lubbe F. // JHEP 2010. V.10. P.44; arXiv:1006.2388.
- [4] Lechtenfeld O., Nölle C., Popov A.D. // JHEP 2010. V.09. P.074; arXiv:1007.0236.
- [5] Chatzistavrakidis A., Lechtenfeld O., Popov A.D. // to appear.
- [6] Gemmer K.P., Haupt A., Lechtenfeld O., Nölle C., Popov A.D. // to appear.